

Nonlinear absorption in discrete systems

A Spire and J Leon

Physique Mathématique et Théorique, CNRS-UMR5825, Université Montpellier 2,
34095 Montpellier, France

Received 15 June 2004, in final form 3 August 2004

Published 15 September 2004

Online at stacks.iop.org/JPhysA/37/9101

doi:10.1088/0305-4470/37/39/004

Abstract

In the context of nonlinear scattering, a continuous wave incident onto a nonlinear discrete molecular chain of coupled oscillators can be partially absorbed as a result of a three-wave resonant interaction that couples two HF-waves of frequencies close to the edge of the Brillouin zone. Hence both nonlinearity and discreteness are necessary for generating this new absorption process which manifests itself by soliton generation in the medium. As a paradigm of this *nonlinear absorption* we consider here the Davydov model that describes exciton–phonon coupling in hydrogen-bonded molecular chains.

PACS numbers: 02.60.Cb, 05.45.–a, 45.05.+x

1. Introduction

The scattering of waves becomes extremely rich when nonlinearity comes into play, and one of its most fundamental effects is *soliton generation* which results in energy localization and propagation. For instance incident radiation on a two-level medium at the resonant frequency can be totally transmitted, instead of being absorbed, a property described in [1] as *self-induced transparency*. It results from the nonlinear coupling of radiation with medium population, a mechanism which generates the solitons, vectors of energy transmission.

Other interesting processes of nonlinear wave scattering are two-photon propagation, second harmonic generation and stimulated Raman scattering. In the last case, the nonlinear interaction induces (laser) pump depletion and phase effects result in Raman spike generation (short duration pump depletion) [2]. Although Raman spike generation is not a solitonic effect, here also nonlinearity is fundamental tool [3].

Recently discovered, *nonlinear supratransmission* [4, 5] is a nonlinear scattering process where a totally reflective medium switches to high transmissivity above some threshold because of nonlinearity. The bifurcation has its origin in a nonlinear instability [6] which has a wide field of application in generic nonlinear evolutions [7].

Particularly interesting is the application of this concept to the scattering of a continuous-wave laser beam incident onto a Bragg mirror which switches from total reflection to

transmission by means of gap soliton generation [8, 9]. It has been demonstrated that the switch is a manifestation of nonlinear supratransmission which has in particular allowed the computation of an analytic explicit expression of the intensity flux threshold [10].

We are interested here in the concept of *nonlinear absorption* where a medium, transparent to incident radiation in the linear regime, can become absorbent under the contribution of nonlinearity which is due here to coupling of waves of different nature. Such is the case with the Davydov model [11] describing the coupling of high frequency optical phonon waves to low frequency acoustic waves. A generic field of application of the Davydov model is crystalline acetanilide ($C_6H_5NHCOCH_3$) where the acoustic wave represents the hydrogen bond between adjacent molecules, while the optical phonon wave represents the C=O stretching (amide-I mode). This model has been widely studied in its stationary limit where the hydrogen bond is supposed to have a time response much longer than the high frequency C=O stretching. In that situation the resulting simplified model is the nonlinear Schrödinger equation that has been used to interpret the anomalous IR-absorption band by soliton generation [12].

The Davydov model has been recently re-examined in the context of resonant two-wave interaction where a rigorous multi-scale analysis transforms the discrete model to a continuous integrable partial differential equation which keeps the fundamental wave coupling effect and possesses multi-soliton solutions [13]. This two-wave resonant interaction process takes into account an incident high-frequency optical phonon wave that resonates with the low-frequency acoustic wave as soon as the HF-wave group velocity equals the LF-wave phase velocity. This is the Benney criterion [14] that has revealed its efficiency through numerical simulations of the Davydov model [15].

The two-wave resonant process assumes no backward propagation and thus neglects the multiple reflections due to the periodicity of the crystal. We consider here a three-wave interaction process that couples the incident optical phonon wave both to the acoustic excitation and the reflected optical phonon wave. We shall discover that an incident and a reflected high-frequency wave can cooperate resonantly with a low-frequency acoustic wave thanks to the discrete nature of the basic model.

We will demonstrate that this scattering process allows for absorption of incident radiation by a purely nonlinear effect that generates a three-wave soliton in the medium. The nonlinear effect is effectively an absorption as a part of the incident energy flux is transferred to the medium.

2. Three-wave interaction in the Davydov model

2.1. Basic model

Our starting point is the Davydov model [11] for the eigenstate $a_n(t)$ of the amide-I excitation (the corresponding dynamical variable represents the C=O stretching) coupled to the dynamical variable β_n that represents the longitudinal displacement along the hydrogen-bond chain. It reads

$$i\hbar\dot{a}_n = [E_0 + W + \chi(\beta_{n+1} - \beta_n)]a_n - J(a_{n+1} + a_{n-1}), \quad (1)$$

$$M\ddot{\beta}_n = \kappa(\beta_{n+1} - 2\beta_n + \beta_{n-1}) + \chi(|a_n|^2 - |a_{n-1}|^2), \quad (2)$$

where M is the mass of the peptide group, K is the spring constant of the hydrogen bond, E_0 is the energy of the C=O stretching and W is the total energy of the peptide group displacements. The constant J measures the energy of the dipole–dipole interaction of C=O stretching oscillations and χ is a quadratic force constant for the exciton–phonon nonlinear

coupling. In the case of the Davydov–Scott [11, 12] approach, χ becomes a cubic force constant which indicates how strongly chain distortion will trap localized vibrational energy. An overdot stands for derivation with respect to the physical time T .

Upon defining the dimensionless time $t = JT/\hbar$, and the adimensional quantities

$$\Psi_n = a_n \frac{\hbar\chi}{J\sqrt{JM}} \exp\left[\frac{i}{J}(E_0 + W - 2J)t\right], \quad Q_n = \frac{\chi}{J}(\beta_{n+1} - \beta_n), \quad (3)$$

the system becomes

$$i\partial_t \Psi_n + (\Psi_{n+1} - 2\Psi_n + \Psi_{n-1}) = Q_n \Psi_n, \quad (4)$$

$$\partial_t^2 Q_n - V^2(Q_{n+1} - 2Q_n + Q_{n-1}) = |\Psi_{n+1}|^2 - 2|\Psi_n|^2 + |\Psi_{n-1}|^2, \quad (5)$$

which constitutes our basic example of a discrete nonlinear coupled wave system of equations. Note that we are left with a single constant, the adimensional sound velocity

$$V = \frac{\hbar}{J}v_p, \quad v_p = \sqrt{\frac{\kappa}{M}}, \quad (6)$$

where v_p is the phonon velocity (cells per second), and that the coupling parameter χ has been absorbed in the amplitude of $\psi_n(t)$.

2.2. Multi-scale expansion

Following the multi-scale expansion method [16] we assume a representation of the solution $\{\Psi_n(t), Q_n(t)\}$ as a formal series in terms of a small parameter ϵ where space–time dependences occur at a sequence of slow scales. Since it is not necessary to push the series at arbitrary order, we write explicitly the first relevant terms only as

$$\Psi_n(t) = \epsilon\varphi^{(0)}(n_0, x_1, \dots; t_0, t_1, \dots) + \epsilon^2\varphi^{(1)}(n_0, x_1, \dots; t_0, t_1, \dots) + \dots \quad (7)$$

$$Q_n(t) = \epsilon q^{(0)}(n_0, x_1, \dots; t_0, t_1, \dots) + \epsilon^2 q^{(1)}(n_0, x_1, \dots; t_0, t_1, \dots) + \dots \quad (8)$$

and the difference-differential operators must be understood as

$$\nabla_n^\pm \rightarrow \nabla_{n_0}^\pm + \epsilon\partial_{x_1} + \dots, \quad \partial_t \rightarrow \partial_{t_0} + \epsilon\partial_{t_1} + \dots \quad (9)$$

Hereabove the difference operators are defined as

$$\nabla_n^+ \Psi_n = \Psi_{n+1} - \Psi_n, \quad \nabla_n^- \Psi_n = \Psi_n - \Psi_{n-1}, \quad (10)$$

such that the second-order difference appearing in (4) is

$$\nabla_n^2 \Psi_n = \nabla_n^+ \nabla_n^- \Psi_n = \Psi_{n+1} + \Psi_{n-1} - 2\Psi_n. \quad (11)$$

The system (4), (5) at first order gives the linear equations

$$\begin{aligned} \mathcal{L}_0\varphi^{(0)} &= 0, & \mathcal{L}_0 &= i\partial_{t_0} + \nabla_{n_0}^2, \\ L_0q^{(0)} &= 0, & L_0 &= \partial_{t_0}^2 - V^2\nabla_{n_0}^2. \end{aligned} \quad (12)$$

The next order ϵ^2 eventually reads

$$\begin{aligned} \mathcal{L}_0\varphi^{(1)} &= -i\partial_{t_1}\varphi^{(0)} - (\nabla_{n_0}^+ + \nabla_{n_0}^-)\partial_{x_1}\varphi^{(0)} + q^{(0)}\varphi^{(0)}, \\ L_0q^{(1)} &= 2\partial_{t_0}\partial_{t_1}q^{(0)} + V^2(\nabla_{n_0}^+ + \nabla_{n_0}^-)\partial_{x_1}q^{(0)} + \nabla_{n_0}^2|\varphi^{(0)}|^2. \end{aligned} \quad (13)$$

The method then works as follows. Once an explicit solution of the linear system (12) is selected, we express that the evolution (13) of the first-order correction $\{\varphi_n^{(1)}, q_n^{(1)}\}$ must not produce secular growth. This explicitly furnishes the evolution of the fundamental $\{\varphi_n^{(0)}, q_n^{(0)}\}$ in the slow variables x_1 and t_1 . The choice of the linear solution in the variables n_0, t_0 determines the physical problem under study.

2.3. Selection rules

To describe a three-wave interaction process involving incident and backscattered optical phonon waves, we are led to select in the linear system (12) the solution

$$\varphi^{(0)} = A(x_1, t_1) e^{i(k_1 n_0 - \omega_1 t_0)} + B(x_1, t_1) e^{i(-k_2 n_0 - \omega_2 t_0)}, \quad (14)$$

$$q^{(0)} = g(x_1, t_1) e^{i(K n_0 - \Omega t_0)} + \bar{g}(x_1, t_1) e^{-i(K n_0 - \Omega t_0)}, \quad (15)$$

with wave numbers $k_1 > 0$ and $k_2 > 0$ such as to ensure an optical phonon wave as a superposition of an incoming wave of amplitude A and a reflected wave of amplitude B . The acoustic wave can propagate in both directions, thus K can be of either sign. We have explicitly written the variables (x_1, t_1) in the slowly varying amplitudes A , B and g , but of course they depend on all higher order variables (but *not* on the first order ones n_0 and t_0). The above expression is a solution of (12) for the following dispersion relations:

$$\omega_1 = 2(1 - \cos k_1), \quad \omega_2 = 2(1 - \cos k_2), \quad \Omega = 2V \left| \sin \frac{K}{2} \right|. \quad (16)$$

Note that using the relation (3) and the multi-scale derivative laws, we demonstrate that the first-order envelope β_0 of the longitudinal displacement along the hydrogen-bond chain β_n is related to the envelope $g(x_1, t_1)$ of Q_n by

$$\beta_0(x_1, t_1) = -i \frac{J}{2\chi \sin(K)} g(x_1, t_1) \quad (17)$$

then the behaviour of the envelope $g(x_1, t_1)$ will automatically give the behaviour of $\beta_0(x_1, t_1)$.

The resonant wave interaction results from a selection rule for the wave parameters, obtained by examination of the nonlinear terms that occur in (13), namely

$$q^{(0)} \varphi^{(0)} = Ag e^{i(k_1+K)n_0} e^{-i(\omega_1+\Omega)t_0} + A\bar{g} e^{i(k_1-K)n_0} e^{-i(\omega_1-\Omega)t_0} \\ + Bg e^{i(-k_2+K)n_0} e^{-i(\omega_2+\Omega)t_0} + B\bar{g} e^{i(-k_2-K)n_0} e^{-i(\omega_2-\Omega)t_0}.$$

Such terms will combine to either components of $\varphi^{(0)}$ and resonate with corresponding factors in the left-hand side of (13). The evolution of the envelopes will then be obtained by setting to zero the coefficients of resonating terms.

Since the physical context is the resonant interaction of two HF-waves (optical phonon) with a LF acoustic wave, it implies a small value of K (near the centre of the Brillouin zone) and large values of k_1 and k_2 , that is to say close to, but less than, the value π in order to achieve incident and reflected HF-waves. We are then left with the following selection rules. First when the nonlinear terms Ag and $B\bar{g}$ combine respectively with B and A we get

$$k_1 + k_2 = 2\pi - K, \quad K > 0, \quad \omega_1 - \omega_2 = -\Omega. \quad (18)$$

Another possibility is to combine instead $A\bar{g}$ and Bg with B and A to obtain

$$k_1 + k_2 = 2\pi + K, \quad K < 0, \quad \omega_1 - \omega_2 = \Omega. \quad (19)$$

We shall study these two cases and demonstrate that only the first one gives an instability that generates solitons, displayed in figure 1.

It is worth remarking that such a scattering process (involving 2π) is allowed for by the presence of an exponential in the discrete variable (n_0), in other words by the discrete nature of the medium.

Inserting the dispersion relations (16) in equation (18), we obtain after some algebra the following solution:

$$2 \sin(k_1 + K/2) = V, \quad 2 \sin(k_2 + K/2) = -V. \quad (20)$$




Figure 1. Representation of the selection rules (18) ($V = 0.5$) and graph of solution (20).

These equations determine completely the wave numbers k_2 and K from the data of V (physics) and k_1 (incident wave), as soon as one assumes that k_1 and k_2 are close to and less than π . The relations (20) can be expressed in terms of molecular chain parameters (using the dispersion relations (16) and the rescaling time defined before) and give the set of optimal properties to display nonlinear absorption.

Note that this is enough to give the physical parameters V (related to the mass, the dipole interaction constant and force constant) and K which is directly related to the frequency of the phonon mode involved in the coupling process. We then obtain the high frequency modes which could interact to obtain nonlinear absorption. In another way, we can impose the frequency of excitonic modes and look for parameters of the chain needed to obtain a resonant process.

Remark. A particular case is obtained in the limit $K \rightarrow 0$ for which $k_2 \rightarrow -k_1$ and $2 \sin(k_1) = V$. This is the *two-wave* resonant interaction studied in [13] for which a different multi-scale analysis has to be employed (resonant process occurs at larger scales).

2.4. Three-wave resonant interaction

Corresponding to the first selection rule (18), the evolution equations for the envelopes that ensure vanishing of the resonant terms in the evolution (13) read

$$\begin{aligned}
 [\partial_t + 2 \sin k_1 \partial_x]A &= -iB\bar{g}, \\
 [\partial_t - 2 \sin k_2 \partial_x]B &= -iAg, \\
 \left[\partial_t - V^2 \frac{\sin K}{\Omega} \partial_x \right] g &= i \frac{\Omega}{2V^2} B\bar{A}.
 \end{aligned}
 \tag{21}$$

From now on we rename x_1 by x and t_1 by t . The same procedure applied to the second selection rule (19) eventually furnishes

$$\begin{aligned}
 [\partial_t + 2 \sin k_1 \partial_x]A &= -iBg, \\
 [\partial_t - 2 \sin k_2 \partial_x]B &= -iA\bar{g} \\
 \left[\partial_t - V^2 \frac{\sin K}{\Omega} \partial_x \right] g &= i \frac{\Omega}{2V^2} A\bar{B}.
 \end{aligned}
 \tag{22}$$

These are two standard three-wave interaction nonlinear evolutions which are integrable systems on the infinite line $x \in \mathbb{R}$ when Cauchy initial data are prescribed in a space of functions vanishing (together with all their derivatives) at $x \rightarrow \pm\infty$ [17].

Such an initial-boundary value problem is not the one we are interested in as indeed we study the scattering of an incident HF wave of envelope $A(x, t)$ having a prescribed value for all time t at the origin of the medium, i.e. at $x = 0$. Moreover, as $B(x, t)$ stands for the envelope of the reflected wave, its value will be prescribed to vanish at the output $x = L$ for all time. Thus we cannot make use of the inverse scattering transform, unless first reformulated for a boundary-value problem on the finite interval, which is still an open problem.

Note that the system (22), by renaming g as \bar{g} , maps to the first one except for the sign of the inhomogeneous term of the last equation. This change of sign is fundamental as it switches from instability to stability, as described below.

2.5. Stability properties

The problem we consider is thus the scattering of an incident HF wave of envelope $A(x, t)$, belonging to a carrier wave of frequency ω_1 , that generates a backscattered wave of envelope $B(x, t)$ and LF acoustic wave of envelope $g(x, t)$ out of initial vacuum. Thus we perform a linear stability analysis of both systems about the solution

$$A(x, t) = A_c, \quad B = 0, \quad g = 0 \tag{23}$$

with constant A_c corresponding to a continuous wave (CW) irradiation. Mathematically speaking this is an exact solution of both systems (21) and (22) and thus only an instability could produce an effective scattering. Let us seek now a solution as the perturbation

$$A = A_c + \epsilon a e^{-i\nu t}, \quad B = \epsilon b e^{-i\nu t}, \quad g = \epsilon q e^{-i\nu t}. \tag{24}$$

The system (21) at order ϵ then gives the linear system

$$\begin{pmatrix} \nu & 0 & 0 \\ 0 & \nu & A_c \\ 0 & -\bar{A}_c \frac{\Omega}{2V^2} & \nu \end{pmatrix} \begin{pmatrix} a \\ b \\ q \end{pmatrix} = 0 \tag{25}$$

possessing the three eigenvalues $\nu = 0$ and $\nu = \pm i(|A_c|/V)\sqrt{\Omega/2}$. The system is thus unstable ($\Omega > 0$), while the same analysis with system (22) and $g = \epsilon q e^{i\nu t}$ furnishes the real eigenvalues 0 and $\pm(|A_c|/V)\sqrt{\Omega/2}$, and thus stability.

Consequently the linear stability analysis predicts that the selection rule (18) will produce an effective scattering for an incident CW wave. Our purpose now is to demonstrate by numerical simulations that this instability is a *soliton generator* which induces an effective energy absorption of the incident radiation.

3. Numerical simulations

In order to understand the effect of the instability of the solution (23) in system (21), and compare it to system (22), we perform here numerical simulations of those systems under the following initial-boundary value data on the interval $x \in [0, L]$:

$$A(0, t) = A_c, \quad B(L, t) = 0, \quad g(x, 0) = 0. \tag{26}$$

As an illustration figure 2 shows the energy density profile $|A(x, t)|^2$ of the incident wave as a function of x at time $t = 20$ for $A_c = 0.7$. The dashed line represents the acoustic phonon $|g(x, t)|^2$ that has gained the energy lost by the incident optical phonon.




Figure 2. Energy density at time $t = 20$ for optical phonon wave $|A(x, t)|^2$ (solid line) and $|g(x, t)|^2$ (crosses).




Figure 3. Energy of the optical phonon wave $I(t)$ for $A_c = 0.7$ and the values of (29).

This process corresponds to a brutal transfer of energy from the incident wave (which has already settled in the medium) to the medium, as shown in figure 3. The normalized energies carried by the waves of envelopes A and g are defined by

$$I(t) = \frac{1}{L|A_c^2|^2} \int_0^L dx |A(x, t)|^2, \tag{27}$$

$$P(t) = \frac{1}{L|A_c^2|^2} \int_0^L dx |g(x, t)|^2. \tag{28}$$

These are the quantities evaluated in a numerical simulation in figure 3 for the following parameters:

$$\begin{aligned} \omega_1 &= 3.94, & k_1 &= 2.89, \\ \omega_2 &= 3.98, & k_2 &= 2.99, \\ V &= 0.1, & \Omega &= 0.04, & K &= 0.4. \end{aligned} \tag{29}$$

Performing the same numerical simulations in the case of system (22) has never produced any nonlinear energy transfer, confirming the predictions of the linear stability analysis.

4. Conclusion

We have shown that a molecular chain allowing for a wave coupling process, like the Davydov model, can present *nonlinear energy absorption* by resonant interaction, coupling HF to LF waves, with selection rules (18) that are allowed only in discrete systems.

The resulting governing equation, though integrable (for a Cauchy initial value problem on the infinite line) leads to an instability when driven by boundary data on the finite interval. This instability is then the source of soliton formation in the medium, and energy transfers from the incident radiation to the medium excitation.

This process is clearly illustrated by numerical simulations and opens the way to further studies in different contexts where the discreteness is known *a priori* to play a fundamental role.

References

- [1] Mc Call S L and Hahn E L 1969 *Phys. Rev.* **183** 457
- [2] Drühl K, Wenzel R G and Carlsten J L 1983 *Phys. Rev. Lett.* **51** 1171
- [3] Claude C and Leon J 1995 *Phys. Rev. Lett.* **74** 3479
Leon J and Mikhailov A V 1999 *Phys. Lett. A* **253** 33
- [4] Caputo J-G, Leon J and Spire A 2001 *Phys. Lett. A* **283** 129
- [5] Geniet F and Leon J 2002 *Phys. Rev. Lett.* **89** 134102
- [6] Leon J 2003 *Phys. Lett. A* **319** 130
- [7] Geniet F and Leon J 2003 *J. Phys.: Condens. Matter* **15** 2933
- [8] Yariv A 1991 *Optical Electronics* 4th edn (Orlando, FL: Saunders)
- [9] Sankey N D, Prelewitz D F and Brown T G 1992 *Appl. Phys. Lett.* **60** 1427
- [10] Leon J and Spire A 2004 *Phys. Lett. A* **327** 474–80
- [11] Davydov A S 1985 *Solitons in Molecular Systems* (Dordrecht: Reidel)
- [12] Scott A C 1992 *Phys. Rep.* **217** 1
- [13] Boiti M, Leon J, Pempinelli F and Spire A 2004 *J. Phys. A: Math. Gen.* **37** 4243
- [14] Benney D J 1977 *Stud. Appl. Math.* **56** 81
- [15] Kopidakis G, Soukoulis C M and Economou E N 1995 *Phys. Rev. B* **51** 15038
- [16] Whitham G B 1974 *Linear and Nonlinear Waves* (New York: Wiley-Interscience)
- [17] Kaup D J 1976 *Stud. Appl. Math.* **55** 9